



HCEO WORKING PAPER SERIES

Working Paper



HUMAN CAPITAL AND
ECONOMIC OPPORTUNITY
GLOBAL WORKING GROUP

The University of Chicago
1126 E. 59th Street Box 107
Chicago IL 60637

www.hceconomics.org

Opioids and Post-COVID Labor-Force Participation^{*}

Francesco Chiocchio[†]

CEMFI

Jeremy Greenwood[‡]

University of Pennsylvania

Nezih Guner[§]

CEMFI, Banco de España

Karen A. Kopeccky[¶]

FRB Cleveland, Emory University

July 16, 2025

Abstract

At the onset of COVID-19, U.S. labor-force participation fell by about 3 percentage points and remained below pre-pandemic levels three years later. Recovery was slower in states hit harder by the pre-pandemic opioid crisis, measured by age-adjusted overdose death rates. An event study shows that a one-standard-deviation increase in pre-COVID opioid deaths led to a 0.9 percentage point drop in post-COVID labor participation. This effect wasn't due to differences in overall health across states and was stronger among those without a college degree. In high-opioid states, slower recovery was linked to more people leaving the workforce due to disability.

Keywords: Labor-Force Participation, Opioids, COVID-19, Health, Disability

JEL Codes: I12, I14, J11, J12, J21

^{*}Comments by Efi Adamopoulou, Dmitry Arkhangelsky, Pedro Mira, and Tom Zohar, and by seminar participants at the Federal Reserve Bank of Cleveland and CEMFI are greatly appreciated. Francesco Chiocchio gratefully acknowledges financial support from Grant PRE2021-098315 funded by MCIN/AEI/10.13039/501100011033 and by “ESF +”, the Maria de Maeztu Unit of Excellence CEMFI MDM-2016-0684, funded by MCIN/AEI/10.13039/501100011033, and CEMFI. The views expressed herein are those of the authors and not necessarily those of Banco de España, the Eurosystem, the Federal Reserve Bank of Cleveland, or the Federal Reserve System.

[†]Email: <francesco.chiocchio@cemfi.edu.es>

[‡]Email: <do-not-use@jeremygreenwood.net>


[§]Email: <ngunermail@gmail.com>

[¶]Email: <karen.kopeccky@clev.frb.org>

1 Introduction

The COVID-19 pandemic caused a sharp drop in US labor-force participation, and the recovery was slow. As Figure 1 shows, the labor-force participation among prime-age workers (25–54) declined by about 3 percentage points. The decline was more pronounced for those without a college degree. Furthermore, the pace of recovery of the labor supply was significantly different across US states. In December 2022, for example, LFP among prime-age workers was still 5 percentage points below the trend in Michigan, while the LFP gap from the trend disappeared in Virginia. On average, LFP was around 2 percentage points below the trend in December 2022, with a standard deviation of 2 percentage points across US states.

Figure 1: Labor-Force Participation Rate, Ages 25-54



Notes: *Monthly labor-force participation rates among 25-54-year-olds by education. Source: Current Population Survey (CPS).*

The COVID-19 pandemic occurred in the US as another epidemic, the opioid crisis, was unfolding (Cutler and Glaeser, 2021; Alpert et al., 2022; Greenwood, Guner, and Kopecky, 2024). Between 2000 and 2019, nearly half a million people died from an opioid overdose. Deaths from opioid overdoses increased significantly during COVID-19, as shown in Figure 2. The number of deaths rose from around 50,000 in 2019 to more than 80,000 in 2021 and 2022. Deaths from other illicit substances, mostly methamphetamine, also increased during COVID-19. The total number of drug overdose deaths increased from about 70,000 in 2019

to over 106,000 in 2021 and 2022. The increase in drug overdose deaths during this period is also documented by [Mulligan \(2022\)](#) and [Greenwood, Guner, and Kopecky \(2022\)](#).

Figure 2: Monthly Deaths from Overdose and Opioid Overdose



Notes: *Monthly deaths due to overdoses and opioid-related overdoses. Source: CDC-Multiple Cause of Death data.*

Individuals who misuse opioids or other illicit substances are significantly less likely to participate in the labor force compared to those who do not use opioids at all or use them strictly as prescribed ([Greenwood, Guner, and Kopecky, 2022](#)). It is shown here that labor supply recovery after the COVID-19 pandemic was slower in US states with higher pre-pandemic exposure to the opioid crisis using an event study approach. The identification strategy utilizes a treatment effect influenced by an initial moderating condition: the onset of COVID-19 serves as the treatment and pre-COVID opioid exposure acts as the moderator. The central hypothesis posits that while COVID-19 led to increased use of illicit substances nationwide, states with higher pre-existing opioid exposure experienced greater increases, further delaying their labor-force recovery.

Pre-COVID opioid exposure is measured using age-adjusted opioid overdose death rates from 2017, serving as a comprehensive indicator of the opioid crisis's intensity in each state. Additionally, this measure suffers less from measurement error than other measures. The analysis shows that a one-standard-deviation increase in pre-COVID opioid exposure reduces labor-force participation by approximately 0.9 percentage points below the trend following the COVID-19 shock. For instance, moving from a state at the 25th percentile of opioid

exposure, such as Minnesota, to one at the 75th percentile, like Michigan or Pennsylvania, corresponds to a 1.1 percentage point decline in labor-force participation. This is substantial, given that the standard deviation in prime-age labor-force participation throughout the post-COVID period across US states was 3.4 percentage points.

We show that this finding is not driven by general health differences across states. States with higher initial opioid exposure may have poorer overall health or weaker healthcare systems, potentially explaining the observed effects. To test this, we conduct a placebo analysis, replacing opioid exposure with age-adjusted death rates from leading non-opioid-related causes. The results show no significant differences in post-COVID labor-force participation between states with higher and lower non-opioid-related death rates.

Furthermore, we find that the effects vary by demographics. While the results are similar for both men and women, differences emerge by educational attainment and age. The effects are significant for labor-force participation among non-college-educated people, who tend to have higher rates of opioid use, but not for the college-educated. Moreover, the impact of opioids on labor-force participation is more pronounced among individuals aged 44 to 54. The results are robust to alternative measures of pre-COVID opioid exposure, including the fraction of individuals with an opioid use disorder and shipments of medications for opioid addiction treatment. An alternative synthetic control group approach, as outlined in [Abadie and Gardeazabal \(2003\)](#), also yields comparable results.

Finally, we explore a possible mechanism to help interpret the results. Individuals with substance-use disorder typically have lower labor-force participation, are more likely to leave the labor force due to disability, and experience poorer health. During COVID-19, the disparities in labor-force participation and disability rates between nonusers and individuals with substance-use disorders widened. Empirical analysis reveals that states with higher pre-COVID opioid exposure saw a greater post-COVID increase in the share of individuals exiting the labor force due to disability, highlighting the critical role of health-related factors in the sluggish labor supply recovery.

Related Literature This study adds to the recent literature on the opioid epidemic’s impact on labor market outcomes. The link between worsening labor market conditions and increased opioid use has been highlighted by [Hollingsworth, Ruhm, and Simon \(2017\)](#), [Carpenter, McClellan, and Rees \(2017\)](#), [Pierce and Schott \(2020\)](#), and [Venkataramani et al. \(2020\)](#). Others, such as [Krueger \(2017\)](#), [Harris et al. \(2020\)](#), [Powell \(2022\)](#), and [Aliprantis,](#)

[Fee, and Schweitzer \(2023\)](#), have examined how opioid use reduces labor-force participation and employment by leveraging geographic variations in opioid exposure. This paper adds to the literature by examining how pre-existing opioid exposure shaped labor-force participation in response to a large health shock.

Within this literature, exogenous changes in prescription drug formulas or their availability have been explored to tease out the causal effects of opioid use on labor market outcomes. The introduction of an abuse-deterrent version of OxyContin in 2010 is used by [Alpert, Powell, and Pacula \(2018\)](#) to show that it led to a higher number of heroin deaths in states with higher initial OxyContin exposure. The same strategy is also used by [Cho et al. \(2021\)](#) to estimate the negative effects of heroin use on employment and labor-force participation. [Beheshti \(2023\)](#) performs a similar analysis, using as an exogenous shock the change in regulations that made the prescription of hydrocodone more difficult in 2014. Like [Alpert, Powell, and Pacula \(2018\)](#), he compares units affected differently by a treatment (changes in drug availability) and shows that areas that experienced larger reductions in hydrocodone prescriptions experienced relative improvements in labor-force participation and employment.

The findings are also related to the literature on the effects of the COVID-19 pandemic on the labor market. Several studies focused on differences across demographic or socioeconomic characteristics, occupations, and industries in their suitability to remote work and, as a result, on how they are impacted by the epidemic. [Alon et al. \(2020\)](#) and [Albanesi and Kim \(2021\)](#) focus on gender, while [Bartik et al. \(2020\)](#), [Dingel and Neiman \(2020\)](#), [Adams-Prassl et al. \(2022b\)](#), and [Mongey, Pilossoph, and Weinberg \(2021\)](#) highlight the role of occupations and their task contents. Another strand of the literature, which is more closely related to the current analysis, documents labor market dynamics during and after the epidemic. The labor market after the epidemic has been surprisingly tight, with low unemployment and labor-force participation rates ([Coibion, Gorodnichenko, and Weber, 2020](#); [Forsythe et al., 2022](#)). Quits and the number of workers looking for new jobs also increased ([Gittleman, 2022](#); [Barlevy et al., 2024](#)). There has also been a decline in the desired work hours that persisted through the end of 2021, as shown in [Faberman, Mueller, and Şahin \(2022\)](#). [Bagga et al. \(2023\)](#) suggest that the post-COVID period was characterized by a shift in workers' valuation of specific job amenities, mainly remote work, which led to persistent labor reallocation. The current analysis contributes to the literature by focusing on opioid use, a factor influencing labor supply behavior that was significantly impacted during the COVID-19 pandemic.

The rest of the paper is structured as follows: Section 2 discusses the data. Section 3 presents the empirical methodology. Section 4 shows the results from the main specification. Robustness checks are presented in Section 5, while Section 6 highlights potential mechanisms. Section 7 concludes.

2 Data and Motivating Evidence

The empirical analysis is conducted at the US state level, covering each month from January 2018 to January 2023. The primary outcome variable is the labor-force participation rate of prime-age civilians (ages 25 to 54), calculated for each state and month using data from the Current Population Survey (CPS). CPS data on gender and educational attainment are also used to analyze outcomes of different socioeconomic groups. Respondents are classified as college-educated if they have attained at least a bachelor’s degree.

The primary measure of pre-COVID opioid exposure is the age-adjusted mortality rate from opioid overdoses in 2017, sourced from the Centers for Disease Control’s *Multiple Cause of Death* database (CDC-MCOD).¹ An opioid overdose death is identified when the underlying cause of death is a drug overdose, and opioids are listed among the multiple causes. For underlying causes, the following ICD-10 codes are included: X40-X44 (accidental drug poisonings), X60-X64 (intentional self-poisoning by drugs), X85 (assault by drug poisoning), and Y10-Y14 (drug poisoning of undetermined intent). Opioids as a multiple cause of death are identified using codes T40.0 (Opium), T40.1 (Heroin), T40.2 (Other opioids), T40.3 (Methadone), T40.4 (Other synthetic narcotics), and T40.6 (Other and unspecified narcotics).

Opioid-related deaths are used as a comprehensive measure of opioid exposure, encompassing both legal and illegal opioid consumption and accounting for variations in the types of opioids used, such as OxyContin, Fentanyl, and heroin, each differing significantly in potency. Summary statistics for the age-adjusted opioid overdose mortality rates across US states are presented in Table 1. Notably, the age-adjusted opioid death rate rose by over 50 percent between 2019 and 2021. There is also considerable variation in death rates across states, as shown in Figure A1 in Appendix A.1. In 2017, opioid death rates ranged from

¹Available at <https://wonder.cdc.gov/mcd.html>. Causes of death are coded using ICD-10, the 10th revision of the International Classification of Diseases by the World Health Organization.

approximately 3 deaths per 100,000 people in Nebraska and Hawaii to 39 in Ohio and 50 in West Virginia. Furthermore, this disparity across states has widened since the COVID-19 pandemic, as evidenced by the increase in the standard deviation between 2019 and 2021, from 10 to 13.²

Table 1: Age-Adjusted Opioid Overdoses Death Rate

	Mean	<i>sd</i>	Min	Max
2017	16.26	10.52	3.1	49.6
2019	16.66	10.16	3.5	43.0
2021	26.26	13.77	5.7	77.2
2022	26.82	13.03	5.6	72.5

Notes: *Summary statistics for the age-adjusted opioid-related overdose death rate for 2017, 2019, 2021, and 2022. Death rates are computed per 100,000 people. Source: CDC-MCOD.*

The relationship between pre-COVID opioid exposure and post-COVID labor supply recovery across US states is presented in Figure 3. As a measure of recovery, the gap between the observed and predicted labor-force participation for prime-age individuals during the last six months of 2022 is used.³ The predicted post-COVID values are based on state-level regressions of labor-force participation on a linear time trend.⁴ More than two years after the onset of COVID-19, states with higher initial opioid death rates in 2017, such as West Virginia and Ohio, had labor-force participation rates that were well below the predicted values.


Two alternative measures of pre-COVID opioid exposure are also considered in the analysis. The first measure is the percentage of individuals with opioid use disorder in 2017-2018, derived from the National Survey on Drug Use and Health (NSDUH). This annual survey provides national and state-level data on tobacco, alcohol, illicit drug use (including non-medical prescription drug use), and mental health. Opioid use encompasses prescription pain relievers and heroin, assessed based on usage within the past 12 months. Prescription misuse includes any use not directed by a doctor, such as use without a prescription or in greater amounts or frequency than prescribed. All heroin users are classified as misusers. Misusers are further screened for opioid use disorder, identified by health issues, disabil-

²Further details on the demographic and geographic variation in fatal drug overdoses over time can be found in Monnat (2022).

³This period was chosen to capture differences across states in LFP recovery, away from the immediate impact of the COVID-19 shock.

⁴Further details on the estimation of trends are provided in Appendix A.2.

Figure 3: Relationship Between Opioid Exposure and Prime-Age LFP Gap



Notes: *Correlation between the predicted gap in prime-age LFP and age-adjusted opioid-related overdose death rate for 2017. The predicted gap in prime-age LFP is constructed by subtracting the predicted LFP from the observed one. The average of the predicted gap between July and December 2022 is reported. Appendix A.2 describes the construction of the predicted LFP values. Source: CDC-MCOD and CPS.*

ities, or significant life disruptions caused by recurring use. The NSDUH is also used to identify individuals who suffer from any drug use disorder, including cocaine, hallucinogens, heroin, inhalants, methamphetamine, and psychotherapeutics (pain relievers, tranquilizers, sedatives, stimulants).⁵

The second measure of opioid exposure is the total shipment of medications used to treat opioid addiction in each state, primarily methadone and buprenorphine. Data are sourced from the Drug Enforcement Agency’s (DEA) Automation of Reports and Consolidated Orders System (ARCOS).⁶ The DEA provides the quantity (in grams) of these drugs distributed to each 3-digit zip code area, which is then aggregated at the state level. Morphine milligram equivalents (MME) are used to calculate a per capita MME amount for each state in 2017.⁷ Summary statistics for these alternative exposure measures and their correlations with age-adjusted death rates are presented in Table 2. There are also significant differences in these measures across states, with a high correlation between these metrics and the age-adjusted

⁵State-level data on drug use disorder are obtained from the Restricted-Use Data Analysis System (RDAS) of the Substance Abuse and Mental Health Services Administration, as the NSDUH public file lacks geographic indicators.

⁶Available at <https://www.dea diversion.usdoj.gov/arcos/arcos.html>. For details on opioid-treatment medications, see (Mutter and Duchovny, 2022). Naltrexone is excluded from ARCOS as it is not a scheduled drug.

⁷MME standardizes opioid potency relative to morphine, facilitating cross-opioid comparisons. Conversion is based on Cutler and Glaeser (2021).

death rates.

Table 2: Alternative Opioid Exposure Measures

	Mean	<i>sd</i>	Min	Max	Corr. with Death Rate
Death Rate, 2017	16.26	10.52	3.10	49.60	1.00
Disorder, 2017-18 (%)	0.84	0.30	0.32	1.60	0.41
Treatment MME pc, 2017	925.43	601.09	157.06	2,809.07	0.78

Notes: *Summary statistics for measures of state opioid exposure. The second row shows the percentage of people with an opioid use disorder in 2017-2018, obtained from the NSDUH. The third row shows the quantity of drugs to treat opioid use disorder distributed to the state in 2017, measured as MME per capita, obtained from ARCOS. The last column shows Spearman’s rank correlation between these two measures and the age-adjusted death rate.*

A potential concern with our exposure measure is that opioid exposure in 2017 may reflect pre-COVID labor market conditions, suggesting that both post-COVID recovery differences and opioid exposure could stem from shared pre-COVID factors. As indicated above, the interaction between opioids and labor-force participation can be complex, with effects going both ways (Abraham and Kearney, 2020). As shown in Figure 4a, there is no significant relationship between 2017 opioid exposure, measured by opioid-related deaths, and labor-force participation during the 2017–2019 period across states. Moreover, Figure 4b shows that the prime-age LFP of states with higher opioid exposure grew faster than that of lower exposure between 2015 and 2019.⁸

The empirical analysis also includes control variables reflecting differences in COVID-19 intensity and policy responses across states. State-level COVID-19 cases and deaths are taken from the CDC’s COVID-19 Tracker.⁹ Policy data are obtained from the Oxford Tracker Dataset (Hale et al., 2021); specifically, the Stringency Index (covering school and workplace closures and stay-at-home orders) and the Economic Support Index (summarizing income support policies during COVID-19) are used.¹⁰ The average value of these two indexes and their standard deviations across US states are shown in Figure A5 in Appendix A.4.¹¹ Given


⁸Figures A3 and A4 in Appendix A.3 analyze this relationship using the two alternative measures of opioid exposure. There is no significant correlation between the measures and LFP between 2017 and 2019. Meanwhile, the relationship between the two alternative measures and the LFP growth rate between 2015 and 2019 is positive.

⁹Available at <https://covid.cdc.gov/covid-data-tracker/#datatracker-home>.

¹⁰The Oxford Tracker Dataset categorizes various COVID-19 policies into indexes representing different policy strengths, normalized between 0 and 100.

¹¹Ruhm (2024) finds that if all states had imposed COVID-19 restrictions similar to those used in the 10 most restrictive states, excess deaths would have been 10 percent to 21 percent lower.

Figure 4: Relationship Between Opioid Exposure and Prime Age Labor Force Participation



Notes: *Left: Correlation between the average prime-age LFP between 2017 and 2019 and the age-adjusted opioid-related overdose death rate for 2017. Right: Correlation between prime-age LFP growth between 2015 and 2019 and age-adjusted opioid-related overdose death rate for 2017. Source: CDC-MCOD and CPS.*

the varied impact of COVID-19 across industries, time-varying state-level employment by industry is also included as a control in the empirical analysis, using CPS data to construct these measures.

3 Empirical Strategy

The simultaneous impact of COVID-19 across all states makes a standard difference-in-differences estimator unsuitable for analyzing why some states experienced a slower recovery in labor-force participation. The analysis here uses an approach that allows the COVID-19 shock to have varying effects across states, depending on their pre-COVID opioid exposure. This approach relies on three assumptions: (1) the treatment affects all units (states) simultaneously, (2) the treatment effect depends on initial, predetermined conditions unrelated to the treatment, and (3) these conditions vary across units. The core idea is that states with higher pre-COVID opioid exposure experienced a slower labor force recovery, potentially due to greater availability of illicit drugs or a larger population with opioid use experience, which became more salient post-COVID. Importantly, COVID-19 onset is assumed to be independent of initial opioid exposure levels.

This empirical strategy is used by [Alpert, Powell, and Pacula \(2018\)](#) to study the effects of OxyContin’s reformulation, which made it harder to abuse. The reformulation, an event that affects everyone, is found to lead to greater increases in heroin deaths in states with higher pre-reformulation OxyContin exposure. A similar approach is also used by [Beheshti \(2023\)](#) to assess the impact of hydrocodone rescheduling by the Drug Enforcement Agency (DEA) that made it harder for doctors to prescribe. Areas with higher initial hydrocodone prescriptions are found to have more significant improvements in labor-force participation and employment.

The empirical approach employs a dynamic two-way fixed effects model, with labor-force participation as the dependent variable and the interaction between the COVID-19 outbreak (February 2020) and pre-COVID opioid exposure as the key explanatory variable. Pre-COVID exposure is measured in two ways using 2017 age-adjusted opioid death rates. The first measure is a binary variable categorizing states into "high" or "low" exposure groups based on whether their death rates are above or below the median. This method enables straightforward, difference-in-difference-like comparisons but sacrifices granularity. For example, Virginia, with 15 opioid deaths per 100,000, falls just below the median in the low-exposure group, while Utah, with 16 deaths per 100,000, is just above the median in the high-exposure group alongside states like Ohio and West Virginia, which have much higher death rates (39 and 50 per 100,000, respectively). The second measure treats death rates as a continuous variable, preserving finer distinctions in exposure. To facilitate interpretation, this continuous variable is standardized to a mean of 0 and a standard deviation of 1.

The event study analysis is based on the following regression:

$$Y_{s,t} = \alpha_s + \alpha_t + \sum_{T \neq 2020m2} \beta_T \times \mathbb{1}\{t = T\} \times Op. \text{Exp}_s + \delta X_{s,t} + \varepsilon_{s,t}, \quad (1)$$

where $Y_{s,t}$ is the outcome of interest, LFP in state s at time t , and α_s and α_t are the state and time fixed effects. The variable $Op. \text{Exp}_s$ is the pre-COVID opioid exposure for state s , which can be either binary, taking the value of 1 if pre-COVID opioid exposure is higher than the median exposure and 0 otherwise, or continuous. The variable $X_{s,t}$ represents time-varying controls.

The coefficients of interest in equation (1) are the β_T , with $\beta_{Feb.2020}$ normalized to 0. These coefficients represent changes in the outcome variable $Y_{s,t}$ relative to the month before COVID-19, based on differing levels of prior opioid exposure. For instance, when $Op. \text{Exp}_s$ is

the binary exposure measure, a value of $\beta_{March.2020} = -0.1$ indicates that, compared to low-exposure states, labor-force participation in high-exposure states was 10 percentage points lower in March 2020 after controlling for their fixed effects. For the continuous exposure measure, $\beta_{March.2020}$ captures the difference in LFP associated with a one-unit increase in the standardized pre-COVID exposure measure, equivalent to a one-standard-deviation increase in the 2017 age-adjusted opioid death rate.

The control variables in $X_{s,t}$ include monthly COVID-19 case counts (set to 0 for the pre-COVID period) to account for the severity of the epidemic across states. Additionally, the Stringency and Economic Support indexes, which summarize state-level COVID-19 policies, are included as they likely influence economic activity and labor supply incentives. Lastly, a Bartik-style control variable is added to account for time-varying differences in employment structure across states. Following [Di Maggio and Kermani \(2016\)](#), this control is constructed as

$$B_{st} = \sum_k \phi_{s,k,\tau} \frac{\nu_{-s,k,t} - \nu_{-s,k,t-1}}{\nu_{-s,k,t-1}},$$

where $\nu_{-s,k,t}$ are the national employment shares in industry k at time t computed by excluding the state $-s$. Meanwhile, $\phi_{s,k,\tau}$ is the employment share in industry k , in state s , at fixed time $\tau = 2017$. Hence, while national employment has declined everywhere, states with larger employment in certain sectors, such as tourism, might be more intensely affected.

The specification in equation (1) assumes that states with different initial opioid exposure exhibited parallel trends in labor-force participation before the COVID-19 epidemic. This parallel trends assumption allows post-COVID differences to be interpreted as difference-in-difference outcomes. Consequently, the estimated values of β_T should be close to 0 during the pre-COVID period. However, this assumption may be too strong and could be violated in the data. Following [Beheshti \(2023\)](#) and [Dobkin et al. \(2018\)](#), a more flexible specification, which allows for preexisting differential linear time trends across states, is also considered, given by

$$Y_{s,t} = \tilde{\alpha}_s + \tilde{\alpha}_t + \sum_{T > 2020m2} \theta_T \mathbb{1}\{t = T\} \times Op. Exp_s + \sum_S \phi_S \mathbb{1}\{s = S\} \times t + \tilde{\delta} X_{s,t} + \varepsilon_{s,t}, \quad (2)$$

where the term $\sum_S \phi_S \mathbb{1}\{s = S\} \times t$ is the term that allows for state-specific linear time trends. The identifying assumption no longer requires parallel trends across states but

instead relies on similar deviations from potentially different linear trends in each state. In this specification, the coefficients θ_T are estimated only for the post-COVID period. However, the β_T estimates from equation (1) can be used to derive θ_T for the pre-COVID period (Dobkin et al., 2018). If these constructed pre-COVID estimates are close to 0, the linear trend assumption can be considered a good fit for the data. Following Beheshti (2023), equation (1) is referred to as a non-parametric event study specification, while equation (2) is called a parametric event study specification.

4 Opioid Exposure and Slow Labor Supply Recovery

The empirical strategy relies on the assumption that the COVID-19 shock led to differential increases in illicit drug use, depending on the pre-COVID opioid exposure. More specifically, we expect that states with higher exposure should also exhibit higher opioid death rates following the COVID-19 shock or, more broadly, higher overdose deaths from any drugs. To test this "first-stage" effect, equation (1) is estimated with opioid death rates (or death rates from illicit drugs) as the dependent variable, $Y_{s,t}$, and the continuous age-adjusted opioid death rates as our measure of pre-COVID opioid exposure, $Op. Exp_s$.¹²

The findings, shown in Figure 5, reveal that a one-standard-deviation increase in initial opioid exposure is associated with a 0.4-point rise in the monthly opioid death rate by May 2020 and a 2-point increase in the yearly death rate for 2020. These effects are substantial, representing 35 percent of the monthly and 13 percent of the yearly pre-pandemic death rates. Only the non-parametric event study results are presented, as the pre-COVID coefficients β_T do not indicate the presence of pre-trends. Figure B2 in Appendix B.2 shows the post-COVID increase in overdose deaths from any drugs. The results are very similar: a one-standard-deviation increase in initial opioid exposure is associated with a 0.43-point rise in the monthly drug overdose death rate by May 2020 and a 1.3-point increase in the yearly death rate for 2020.¹³


Another assumption of the empirical analysis is that COVID-19 cases (treatment) were

¹²For results using binary exposure, where states are categorized above or below the median 2017 age-adjusted death rate, see Figure B1 in Appendix B.1.

¹³For drug overdose deaths, the following ICD-10 codes are included: X40-X44 (accidental drug poisonings), X60-X64 (intentional self-poisoning by drugs), X85 (assault by drug poisoning), and Y10-Y14 (drug poisoning of undetermined intent).

uncorrelated with pre-COVID opioid exposure (predetermined condition) across states. Figure B3 in Appendix B.3 reports the estimates obtained by running the non-parametric event study, equation (1), using monthly COVID-19 cases per 100,000 people as the dependent variable and the continuous measure of initial opioid exposure as the independent variable, where none of the estimated β_T coefficients are significant.

Figure 5: Non-Parametric Event Study with Continuous Exposure - Opioid Deaths




Notes: Non-parametric event study coefficients β_T and the 95% confidence interval. Coefficients represent changes in the opioid overdose deaths relative to February 2020 between states with different levels of prior opioid exposure. The top panel uses the monthly crude rate, while the bottom uses the yearly age-adjusted rate. The initial opioid exposure measure has been normalized to have a standard deviation equal to 1.

The results for labor-force participation are presented next. The β_T estimates from the non-parametric event study are shown in Figure 6. To ease interpretation, coefficients are already multiplied by 100 and presented as percentage points. The upper panel presents results from equation (1) using a binary measure of exposure, categorizing states as above or below the median 2017 age-adjusted opioid death rate. The lower panel shows results with the continuous measure of opioid exposure.

In both cases, the estimated effects are primarily negative, suggesting lower labor-force participation post-COVID in states with higher pre-COVID opioid exposure. However, these estimates are generally not statistically significant, except for February and March 2022. Additionally, the consistent negativity of β_T coefficients in the pre-COVID period indicates

the presence of differential pre-trends in prime-age labor-force participation.

Figure 6: Non-Parametric Event Study, Labor-Force Participation



Notes: Non-parametric event study coefficients β_T and the 95% confidence interval. Coefficients represent changes in prime-age LFP relative to February 2020 between states with different levels of prior opioid exposure. For the top panel, the initial opioid exposure measure has been dichotomized: states with an age-adjusted opioid death rate in 2017 above the median are given a value of 1, and the others a value of 0. For the bottom panel, the initial opioid exposure measure has been normalized to have a standard deviation equal to 1.


The estimates from the parametric event study, based on equation (2), which account for state-specific linear trends, are presented in Figure 7. The upper panel displays results using the binary measure of opioid exposure, while the lower panel shows results using the continuous measure. To ease interpretation, coefficients are again multiplied by 100 and presented as percentage points. For both specifications, the pre-COVID estimated coefficients are approximately 0, supporting the identifying assumption that states experienced similar deviations from linear trends. For this reason, this is our preferred specification.

Post-COVID, the estimated θ_T coefficients indicate that labor-force participation rates in states with higher pre-COVID opioid exposure consistently fell below their linear trend compared to less-exposed states. The gap in labor-force participation between states with higher and lower opioid exposure emerged immediately after COVID-19 and continued to widen, reaching nearly 1 percentage point by January 2021 and approximately 2 percentage points by January 2022. The upper panel estimates indicate that from March 2020 to January 2023, LFP in states with above-median opioid exposure averaged 1.1 percentage

points below its trend.

Similarly, the continuous measure in the lower panel shows that a one-standard-deviation increase in 2017 age-adjusted opioid death rates corresponds to an additional 0.9 percentage point drop in LFP during the post-COVID period. This implies that moving from the 25th percentile of initial opioid exposure (e.g., Minnesota) to the 75th percentile (e.g., Pennsylvania or Michigan) leads to an average LFP decline of 1.1 percentage points—a substantial effect, given that the standard deviation in prime-age LFP across states during the post-COVID period was 3.4 percentage points.

Figure 7: Parametric Event Study, Labor-Force Participation



Notes: *Parametric event study coefficients θ_T and the 95% confidence interval. Coefficients represent average differences from state trends in prime-age LFP between states with different levels of prior opioid exposure. Top panel, the initial opioid exposure measure has been dichotomized: states with an age-adjusted opioid death rate in 2017 above the median are given a value of 1, and the others a value of 0. For the bottom panel, the initial opioid exposure measure has been normalized to have a standard deviation equal to 1.*

4.1 The Effect of Overall Health

A potential concern with the analysis is that states with higher initial opioid exposure may also have weaker healthcare systems or poorer overall health. If this were the case, it could imply that the slower LFP recovery in more exposed states is not associated with

higher exposure to the opioid epidemic but simply with worse overall health.

To address this concern, we conducted a placebo analysis by replacing initial pre-COVID opioid exposure in the main analysis with age-adjusted death rates from non-opioid-related causes. If these placebo results are significant, it would suggest that the larger drop in labor-force participation in states with higher opioid exposure may be attributed to broader health problems in the population or state-specific characteristics of the healthcare sector that influence both pre-COVID opioid mortality and post-COVID labor supply decisions.

For the placebo analysis, we use data from the CDC Underlying Cause of Death dataset on the 15 leading causes of death in the United States.¹⁴ Table B1 in Appendix B.4 lists these causes along with their age-adjusted death rates for 2017. Deaths due to accidents and intentional self-harm (causes 3 and 8) may overlap with opioid-related deaths, and hence are excluded.¹⁵ Figure 8 presents the estimated coefficients from the parametric event study where the initial exposure measure is based on the total death rate from the remaining 13 causes.¹⁶ The coefficients for the post-COVID period are not statistically significant.¹⁷

4.2 Heterogeneity

The results for different demographic and socioeconomic groups are presented using the parametric event study with a continuous exposure measure, as specified in equation (2). Effects across gender and age groups are shown in Figure 9. The estimates are similar for men and women but show stronger effects among older individuals (ages 45–54) compared to younger ones (ages 25–44). Specifically, a one-standard-deviation increase in pre-COVID opioid exposure is associated with a 1.3 percentage point decline in labor-force participation for the 45–54 age group, compared to 0.7 percentage point for the 25–44 age group. Results for age groups 18–24 and 55–64, which are not statistically significant, are shown in Figure B6 in Appendix B.5.¹⁸

¹⁴<https://wonder.cdc.gov/ucd-icd10.html>.


¹⁵Accidents include underlying causes with ICD-10 codes X40-X44, and intentional self-harm includes ICD-10 codes X60-X64.

¹⁶Using the non-parametric event study specification leads to similar results.

¹⁷Results using the top 10 leading causes of death excluding accidents and intentional self-harm deaths, and each of the 13 underlying causes as the initial exposure measure, are shown in Figures B4 and B5 in Appendix B.4.

¹⁸In principle, the analysis in the previous section can be replicated at the county level. The main source of data, the Bureau of Labor Statistics Local Area Unemployment Statistics (LAUS), however, only provides

Figure 8: Parametric Event Study with Placebo Exposure



Notes: *Placebo parametric event study coefficients θ_T and the 95% confidence interval when using prime-age LFP as dependent. Coefficients represent average differences from state trends in prime-age LFP between states with different levels of prior placebo exposure. The measure of prior placebo exposure is the total age-adjusted death rate in 2017 due to the top 15 underlying causes of death. The death rate has been normalized to have a standard deviation equal to 1.*

Effects by educational attainment are illustrated in Figure 10 and show notable differences. For non-college-educated individuals, a one-standard-deviation increase in pre-COVID opioid exposure corresponds to a 1.1 percentage point decline in LFP relative to the linear trend. In contrast, the estimates for college graduates are not significantly different from zero. This result is in line with evidence that labor market outcomes for non-college-educated individuals are more affected by opioids (Aliprantis, Fee, and Schweitzer, 2023).


5 Robustness

Several sensitivity and robustness checks are conducted. First, alternative measures of pre-COVID opioid exposure are used to validate the results obtained with our preferred

an aggregate measure of the labor force. Hence, it is not possible to focus the analysis on prime-age workers. The results, while consistent with state-level analysis, are less precise, which is not surprising given our findings for different age groups.

Figure 9: Parametric Event Study with Continuous Exposure

Labor-Force Participation by Gender and Age



Notes: *Parametric event study coefficients θ_T and the 95% confidence interval. Coefficients represent average differences from state trends in prime-age LFP between states with different levels of prior opioid exposure. The top panel divides the sample by gender, displaying the prime-age labor force participation (LFP) among males and females. The bottom panels show LFP by age groups: 25-44 in the left panel and 45-54 in the right. The initial opioid exposure measure has been normalized to have a standard deviation equal to 1.*


specification, i.e., the parametric event study given in equation (2). Second, a synthetic control method is applied as an alternative empirical approach.

Alternative Exposure Measures: The analysis uses age-adjusted opioid death rates from 2017 as a measure of opioid exposure. First, the single-year measure is replaced with the average age-adjusted death rates from 2010 to 2017. Second, given the slightly right-skewed distribution of 2017 death rates across states, estimates are provided using the logarithm of death rates as the exposure measure. Figure B7 in Appendix B.6 presents the results using these variants of pre-COVID opioid exposure within the parametric event study specification with continuous exposure. The post-COVID gap in labor-force participation closely aligns with the estimates in Figure 7.

Next, results using two alternative measures of opioid exposure are presented. The first is the percentage of individuals aged 12 and above with an opioid use disorder, as

Figure 10: Parametric Event Study with Continuous Exposure

Labor-Force Participation by Education




Notes: *Parametric event study coefficients θ_T and the 95% confidence interval. Coefficients represent average differences from state trends in prime-age LFP between states with different levels of prior opioid exposure. The upper panel focuses on prime-age LFP for non-college-educated individuals, while the bottom panel focuses on college-educated individuals. The initial opioid exposure measure has been normalized to have a standard deviation equal to 1.*

reported in the NSDUH. The second is the per-capita shipment of medications for opioid addiction treatment, sourced from ARCOS. Figure 11 shows estimates from a parametric event study with the continuous exposure measure. Despite a smaller magnitude, the results closely mirror those in Figure 7. States with either a one-standard-deviation higher share of individuals with an opioid use disorder or a one-standard-deviation higher per-capita supply of opioid treatment medications experienced about a 0.6 percentage point larger negative deviation from the trend in labor-force participation.

Alternative Empirical Approach: The parametric event study approach assumes that states with different initial opioid exposure experienced similar deviations from their state-specific linear trends. As an alternative empirical strategy, results using a *synthetic control* method, following Abadie and Gardeazabal (2003) and Abadie, Diamond, and Hainmueller (2010), are presented.

In this method, states are classified as treated or untreated based on whether their pre-COVID opioid exposure is above or below the median. Weights are assigned to states in

Figure 11: Parametric Event Study - Alternative Measures of Opioid Exposure




Notes: *Parametric event study coefficients θ_T and the 95% confidence interval. Coefficients represent average differences from a state trend in prime-age LFP between states with different levels of prior opioid exposure. For prior opioid exposure, the top panel uses the percentage of people with a substance use disorder in 2017-2018. The bottom panel uses the amount of MME per capita of drugs to treat opioid use disorder distributed to the state in 2017. The initial opioid exposure measure has been normalized to have a standard deviation equal to 1.*

the untreated group to construct a synthetic control group that closely matches the average pre-COVID labor-force participation of the treated group. These weights are time-invariant and are determined by minimizing the difference in pre-COVID LFP between the synthetic control and treated groups. By design, the synthetic control group mimics the pre-COVID trend in LFP of the treated group. The treatment effect is then estimated by comparing post-COVID LFP between the treated group and the synthetic control group, calculated as the difference in average LFP during the post-COVID period.

Figure 12 illustrates the average prime-age labor-force participation in treated states and the synthetic control group. In the post-COVID period, LFP diverged between the two groups, with treated states—those with higher opioid exposure—exhibiting lower LFP. The average treatment effect in the post-COVID period is -0.7 percentage points, and it is statistically significant at the 5 percent level.¹⁹

¹⁹Significance is based on bootstrapped standard errors for the average treatment effect over the period, with 5000 repetitions.

Figure 12: Synthetic Control and Treatment Group - LFPR



Notes: *Labor-force participation in treatment and synthetic control groups. The treatment group includes states with an initial opioid exposure above the median. The LFP of the synthetic control is obtained as a weighted average of the LFP of states with an initial opioid exposure below the median.*

6 Mechanisms

The COVID-19 pandemic coincided with an unprecedented rise in deaths from opioids and other illicit drugs, driven by multiple factors. Pandemic-induced isolation worsened mental health (Panchal et al., 2020; Adams-Prassl et al., 2022a), while disruptions in medical services reduced access to medications for opioid use disorder (Currie et al., 2021a; Russell et al., 2021). This period also saw a shift in opioid consumption toward fentanyl, a more potent opioid (Currie et al., 2021b). Additionally, increased free time, lower drug prices, and government income support may have exacerbated the use of illicit substances (Mulligan, 2022).

Although drug overdose deaths rose nationwide during COVID-19, the impact was most severe in states with higher pre-COVID opioid exposure, as shown in Figures 5 and B2. This differential effect may reflect established illegal opioid distribution networks, more lenient prescribing practices, or a larger population experienced with opioids in highly affected states.

The greater increase in illicit drug use, proxied by opioid overdose deaths in states

with higher initial opioid exposure, may have contributed to slower labor-force participation recovery in the post-COVID period, as drug users typically have lower participation rates and poorer health. Table 3 highlights these disparities. In 2019, before COVID-19, 17 percent of individuals aged 24–49 who did not misuse any drugs (excluding marijuana) were out of the labor force, compared to 26 percent of those with a drug use disorder.²⁰

Individuals with a drug use disorder are more likely to report disability as the reason they are not participating in the labor market. As Table 3 reports, in 2019, 17 percent of nonparticipants who did not use any drugs reported disability as the reason compared with 25 percent of those with a drug use disorder. Drug misuse also impacts job attendance; those with drug use disorders missed an average of 4.1 workdays per month, compared to 1.1 days among nonusers. Self-reported health outcomes were similarly much poorer: 26 percent of individuals with drug use disorders reported fair or poor health, compared to 10 percent among nonusers. Finally, comparing 2019 data with the 2021–2022 period, Figure 13 shows that the LFP and disability gaps between nonusers and those with drug use disorders widened after COVID-19.²¹

Table 3: Employment and Health by Drug Use, Ages 24-49

<i>Panel A: Employment</i>	Out of LF	Disability	Total Skipped Days
Nonuser	17%	17%	1.1
Disorder	26%	25%	4.1
<i>Panel B: Self-Reported Health</i>	Very Good	Good	Fair/Poor
Nonuser	61%	29%	10%
Disorder	40%	35%	26%


Notes: *Employment-related variables in Panel A, and self-reported health status in Panel B, by type of drug user in 2019. In Panel A, the first column displays the share of individuals who are not in the labor force. The second column shows the share of individuals out of the labor force who report disability as the reason for not participating. The third column displays the average number of workdays missed in the past 30 days due to illness or other reasons. Self-reported health status is a categorical variable with 4 options: very good, good, fair, and poor. Fair and poor categories are aggregated. Nonusers did not use any drug (excluding marijuana) in the past 12 months. Disorder indicates a drug use disorder (excluding marijuana) in the last 12 months. Source: NSDUH.*

If higher rates of drug use were indeed having a greater impact on the labor market in

²⁰This age group is used because NSDUH reports age in brackets.

²¹Tables C1 and C2 in Appendix C.1 provide detailed employment and health outcomes by drug use in 2019 and 2021.

Figure 13: Changes in Labor-Force Participation by Drug Use, Ages 24-49



Notes: *Out of labor force variables by type of drug use in 2019 and 2021–2022. Left panel: share of individuals who are out of the labor force. Right panel: share of individuals out of the labor force who report disability as the reason they are not participating. Nonusers did not use any drug (excluding marijuana) in the past 12 months. Disorder indicates a drug use disorder (excluding marijuana) in the last 12 months. Source: NSDUH.*


states with higher pre-COVID opioid exposure, we should see a larger increase in the share of individuals out of the labor force due to disability in those states. This follows from the correlation between drug use disorder and reporting disability as the reason for nonparticipation. To explore the relationship between pre-COVID opioid exposure and nonparticipation due to disability, information is obtained from the CPS, where individuals not participating in the labor force select a reason. This can be being "retired," being "unable to work" (for medical conditions that prevent work for six months or more), or "other." Those who selected the "other" category may specify a reason, including "disability," "illness," being "in school," or "taking care of house or family."

Using this information, the prime-age out-of-labor-force due to disability rate is defined as the share of prime-age individuals who do not participate in the labor force and have as their reason "unable to work" or "disability." The out-of-labor-force due to disability rate has been on a declining trend since 2015, driven mainly by those without a college degree. The decline, however, stopped with the pandemic, stabilizing at around 0.05 (Figure C1 in Appendix C.2).

To analyze the post-COVID trajectory across states of the share of prime-age individuals

who are not working due to disability, a parametric event study using a continuous measure of initial opioid exposure, as specified in equation (2), is estimated with the out-of-labor-force due to disability rate as the dependent variable. The estimated coefficients are presented in Figure 14. The results indicate that pre-COVID, states with varying opioid exposure levels exhibited similar deviations from their trends. Post-COVID, however, states with higher preexisting opioid exposure deviated more positively from these trends. Specifically, a one-standard-deviation increase in opioid exposure is associated with a 0.4 percentage point rise in the post-COVID out-of-labor-force due to disability rate. This is a substantial effect, given that the standard deviation of the out-of-labor-force due to disability rate during the post-COVID period was 2.0 percentage points.²²

Figure 14: Parametric Event Study Estimates - Out of LF due to Disability



Notes: *Parametric event study coefficients θ_T and the 95% confidence interval. Coefficients represent average differences from state trends in prime-age out-of-labor-force due to disability rates between states with different levels of prior placebo exposure. The out-of-labor-force due to disability rate is defined as the proportion of individuals out of the labor force due to disability to the total population. The initial opioid exposure measure has been normalized to have a standard deviation equal to 1.*

²²As with labor-force participation, these results are primarily driven by the non-college-educated population, as shown in Figure C2 in Appendix C.3.

7 Conclusions

The COVID-19 pandemic significantly disrupted labor-force participation across the United States, with prolonged declines among prime-age individuals, particularly those without a college degree. This study demonstrates that states with higher pre-pandemic opioid exposure experienced slower labor-force recovery, underscoring how public health crises can compound labor market challenges. The findings suggest that preexisting patterns of opioid misuse intensified the labor market disruptions caused by COVID-19, likely due to increased substance abuse driven by pandemic-related factors. The interaction between the COVID-19 and opioid crises highlights that the labor force impacts of future economic shocks may similarly be shaped by preexisting public health vulnerabilities.

References

- Abadie, Alberto, Alexis Diamond, and Jens Hainmueller.** 2010. “Synthetic Control Methods for Comparative Case Studies: Estimating the Effect of California’s Tobacco Control Program.” *Journal of the American Statistical Association*, 105(490): 493–505. <https://doi.org/10.1198/jasa.2009.ap08746>.
- Abadie, Alberto, and Javier Gardeazabal.** 2003. “The Economic Costs of Conflict: A Case Study of the Basque Country.” *American Economic Review*, 93(1): 113–132. <https://doi.org/10.1257/000282803321455188>.
- Abraham, Katharine G, and Melissa S Kearney.** 2020. “Explaining the Decline in the US Employment-to-Population Ratio: A Review of the Evidence.” *Journal of Economic Literature*, 58(3): 585–643. <https://doi.org/10.1257/jel.20191480>.
- Adams-Prassl, Abi, Teodora Boneva, Marta Golin, and Christopher Rauh.** 2022*a*. “The Impact of the Coronavirus Lockdown on Mental Health: Evidence from the United States.” *Economic Policy*, 37(109): 139–155. <https://doi.org/10.1093/epolic/eiac002>.
- Adams-Prassl, Abi, Teodora Boneva, Marta Golin, and Christopher Rauh.** 2022*b*. “Work That Can Be Done from Home: Evidence on Variation within and across Occupations and Industries.” *Labour Economics*, 74: 102083. <https://doi.org/10.1016/j.labeco.2021.102083>.
- Albanesi, Stefania, and Jiyeon Kim.** 2021. “Effects of the COVID-19 Recession on the US Labor Market: Occupation, Family, and Gender.” *Journal of Economic Perspectives*, 35(3): 3–24. <https://doi.org/10.1257/jep.35.3.3>.
- Aliprantis, Dionissi, Kyle Fee, and Mark E. Schweitzer.** 2023. “Opioids and the Labor Market.” *Labour Economics*, 85: 102446. <https://doi.org/10.1016/j.labeco.2023.102446>.
- Alon, Titan, Matthias Doepke, Jane Olmstead-Rumsey, and Michèle Tertilt.** 2020. “This Time It’s Different: The Role of Women’s Employment in a Pandemic Recession.” National Bureau of Economic Research Working Paper 27660. <https://doi.org/10.3386/w27660>.
- Alpert, Abby, David Powell, and Rosalie Liccardo Pacula.** 2018. “Supply-Side Drug Policy in the Presence of Substitutes: Evidence from the Introduction of

- Abuse-Deterrent Opioids.” *American Economic Journal: Economic Policy*, 10(4): 1–35. <https://doi.org/10.1257/pol.20170082>.
- Alpert, Abby, William N. Evans, Ethan M.J. Lieber, and David Powell.** 2022. “Origins of the Opioid Crisis and Its Enduring Impacts.” *The Quarterly Journal of Economics*, 137(2): 1139–1179. <https://doi.org/10.1093/qje/qjab043>.
- Bagga, Sadhika, Lukas Mann, Aysegül Şahin, and Giovanni L. Violante.** 2023. “Job Amenity Shocks and Labor Reallocation.” Working Paper.
- Barlevy, Gadi, R. Jason Faberman, Bart Hobijn, and Aysegül Şahin.** 2024. “The Shifting Reasons for Beveridge Curve Shifts.” *Journal of Economic Perspectives*, 38(2): 83–106. <https://doi.org/10.1257/jep.38.2.83>.
- Bartik, Alexander W., Zoe B. Cullen, Edward L. Glaeser, Michael Luca, and Christopher T. Stanton.** 2020. “What Jobs are Being Done at Home During the Covid-19 Crisis? Evidence from Firm-Level Surveys.” National Bureau of Economic Research Working Paper 27422. <https://doi.org/10.3386/w27422>.
- Beheshti, David.** 2023. “The Impact of Opioids on the Labor Market: Evidence from Drug Rescheduling.” *Journal of Human Resources*, 58(6): 2001–2041. <https://doi.org/10.3368/jhr.0320-10762R1>.
- Carpenter, Christopher S., Chandler B. McClellan, and Daniel I. Rees.** 2017. “Economic Conditions, Illicit Drug Use, and Substance Use Disorders in the United States.” *Journal of Health Economics*, 52: 63–73. <https://doi.org/10.1016/j.jhealeco.2016.12.009>.
- Cho, David, Daniel Garcia, Joshua Montes, and Alison Weingarden.** 2021. “Labor Market Effects of the Oxycodone-Heroin Epidemic.” FEDS Working Paper 25. <https://doi.org/10.17016/FEDS.2021.025>.
- Coibion, Olivier, Yuriy Gorodnichenko, and Michael Weber.** 2020. “Labor Markets During the COVID-19 Crisis: A Preliminary View.” National Bureau of Economic Research Working Paper 27017. <https://doi.org/10.3386/w27017>.
- Currie, Janet M., Molly K. Schnell, Hannes Schwandt, and Jonathan Zhang.** 2021a. “Prescribing of Opioid Analgesics and Buprenorphine for Opioid Use Disorder During the COVID-19 Pandemic.” *JAMA Network Open*, 4(4): e216147–e216147. <https://doi.org/10.1001/jamanetworkopen.2021.6147>.

- Currie, Janet M., Molly K. Schnell, Hannes Schwandt, and Jonathan Zhang. 2021b. “Trends in Drug Overdose Mortality in Ohio During the First 7 Months of the COVID-19 Pandemic.” *JAMA Network Open*, 4(4): e217112–e217112. <https://doi.org/10.1001/jamanetworkopen.2021.7112>.
- Cutler, David M, and Edward L. Glaeser. 2021. “When Innovation Goes Wrong: Technological Regress and the Opioid Epidemic.” *Journal of Economic Perspectives*, 35(4): 171–196. <https://doi.org/10.1257/jep.35.4.171>.
- Di Maggio, Marco, and Amir Kermani. 2016. “The Importance of Unemployment Insurance as an Automatic Stabilizer.” National Bureau of Economic Research. <https://doi.org/10.3386/w22625>.
- Dingel, Jonathan I., and Brent Neiman. 2020. “How Many Jobs Can be Done at Home?” *Journal of Public Economics*, 189: 104235. <https://doi.org/10.1016/j.jpubeco.2020.104235>.
- Dobkin, Carlos, Amy Finkelstein, Raymond Kluender, and Matthew J. Notowidigdo. 2018. “The Economic Consequences of Hospital Admissions.” *American Economic Review*, 108(2): 308–352. <https://doi.org/10.1257/aer.20161038>.
- Faberman, R. Jason, Andreas I. Mueller, and Aysegül Şahin. 2022. “Has the Willingness to Work Fallen during the Covid Pandemic?” *Labour Economics*, 79: 102275. <https://doi.org/10.1016/j.labeco.2022.102275>.
- Forsythe, Eliza, Lisa B. Kahn, Fabian Lange, and David Wiczer. 2022. “Where Have All the Workers Gone? Recalls, Retirements, and Reallocation in the COVID Recovery.” *Labour Economics*, 78: 102251. <https://doi.org/10.1016/j.labeco.2022.102251>.
- Gittleman, Maury. 2022. “The “Great Resignation” in perspective.” *Monthly Labor Review*, U.S. Bureau of Labor Statistics, July. <https://doi.org/10.21916/mlr.2022.20>.
- Greenwood, Jeremy, Nezih Guner, and Karen Kopecky. 2022. “Substance Abuse during the Pandemic: Implications for Labor-Force Participation.” *HSOA Journal of Addiction and Addictive Disorders*, 9(2): 100087. <http://doi.org/10.24966/AAD-7276/100087>.
- Greenwood, Jeremy, Nezih Guner, and Karen Kopecky. 2024. “The Downward Spiral: A Macroeconomic Analysis of the Opioid Crisis.” Federal Reserve Bank of Cleveland Working Paper WP 24-18. <https://doi.org/10.26509/frbc-wp-202418>.

- Hale, Thomas, Noam Angrist, Rafael Goldszmidt, Beatriz Kira, Anna Petherick, Toby Phillips, Samuel Webster, Emily Cameron-Blake, Laura Hallas, Saptarshi Majumdar, et al. 2021. “A Global Panel Database of Pandemic Policies (Oxford COVID-19 Government Response Tracker).” *Nature Human Behaviour*, 5(4): 529–538. <https://doi.org/10.1038/s41562-021-01079-8>.
- Harris, Matthew C., Lawrence M. Kessler, Matthew N. Murray, and Beth Glenn. 2020. “Prescription Opioids and Labor Market Pains: The Effect of Schedule II Opioids on Labor Force Participation and Unemployment.” *Journal of Human Resources*, 55(4): 1319–1364. <https://doi.org/10.3368/jhr.55.4.1017-9093R2>.
- Hollingsworth, Alex, Christopher J Ruhm, and Kosali Simon. 2017. “Macroeconomic Conditions and Opioid Abuse.” *Journal of Health Economics*, 56: 222–233. <https://doi.org/10.1016/j.jhealeco.2017.07.009>.
- Krueger, Alan B. 2017. “Where Have All the Workers Gone? An Inquiry into the Decline of the US Labor Force Participation Rate.” *Brookings Papers on Economic Activity*, 2017(2): 1. <https://doi.org/10.1353/eca.2017.0012>.
- Mongey, Simon, Laura Pilossoph, and Alexander Weinberg. 2021. “Which Workers Bear the Burden of Social Distancing?” *The Journal of Economic Inequality*, 19(3): 509–526. <https://doi.org/10.1007/s10888-021-09487-6>.
- Monnat, Shannon M. 2022. “Demographic and Geographic Variation in Fatal Drug Overdoses in the United States, 1999–2020.” *The ANNALS of the American Academy of Political and Social Science*, 703(1): 50–78. <https://doi.org/10.1177/00027162231154348>.
- Mulligan, Casey B. 2022. “Lethal Unemployment Bonuses? Substitution and Income Effects on Substance Abuse, 2020–21.” National Bureau of Economic Research Working Paper 29719. <https://doi.org/10.3386/w29719>.
- Mutter, Ryan, and Noelia Duchovny. 2022. “Recent Federal Responses to the Opioid Crisis.” *NEJM Catalyst Innovations in Care Delivery*, 3(6).
- Panchal, Nirmita, Rabah Kamal, Kendal Orgera, Cynthia Cox, Rachel Garfield, Liz Hamel, and Priya Chidambaram. 2020. “The Implications of COVID-19 for Mental Health and Substance Use.” *Kaiser Family Foundation*, 21: 1–16.

- Pierce, Justin R, and Peter K Schott.** 2020. “Trade Liberalization and Mortality: Evidence from US Counties.” *American Economic Review: Insights*, 2(1): 47–63. <https://doi.org/10.1257/aeri.20180396>.
- Powell, David.** 2022. “The Labor Supply Consequences of the Opioid Crisis.” RAND Education and Labor Working Paper Series WR-A1484-1. <https://doi.org/10.7249/WRA1484-1>.
- Ruhm, Christopher J.** 2024. “US State Restrictions and Excess COVID-19 Pandemic Deaths.” *JAMA Health Forum*, 5(7): e242006–e242006. <https://doi.org/10.1001/jamahealthforum.2024.2006>.
- Russell, Cayley, Farihah Ali, Frishta Nafeh, Jürgen Rehm, Sean LeBlanc, and Tara Elton-Marshall.** 2021. “Identifying the impacts of the COVID-19 pandemic on service access for people who use drugs (PWUD): A national qualitative study.” *Journal of Substance Abuse Treatment*, 129: 108374. <https://doi.org/10.1016/j.jsat.2021.108374>.
- Venkataramani, Atheendar S., Elizabeth F. Bair, Rourke L. O’Brien, and Alexander C. Tsai.** 2020. “Association Between Automotive Assembly Plant Closures and Opioid Overdose Mortality in the United States: A Difference-in-Differences Analysis.” *JAMA Internal Medicine*, 180(2): 254–262. <https://doi.org/10.1001/jamainternmed.2019.5686>.


Online Appendix

A Data

A.1 Opioid Deaths Across the United States

Figure A1 shows the age-adjusted opioid-related overdose death rates in 2017 across the United States. The regions with the highest death rates, ranging from 16 to 50 per 100,000, are the Appalachian region, the Rust Belt, and New England.

Figure A1: Age-Adjusted Death Rates from Opioid Overdose, 2017



Notes: Age-adjusted opioid-related overdose death rate for 2017 across the US. Death rates are computed per 100,000 people. Source: CDC-MCOD.

A.2 Labor Force Participation Trends

To obtain the difference between the actual and predicted prime-age labor-force participation shown in Figure 3, we do the following steps. First, we estimate a linear time trend in prime-age LFP for each state from January 2010 to February 2020. Second, we use the

estimated coefficients to forecast the prime-age LFP from March 2020 to December 2022. Lastly, we determine the difference between the observed values and the trend.


To predict the trend, we first HP filter the time series for each state for the period January 2010 - February 2020 (default smoothing parameter, 1600×34). Then, we run on the obtained trend the following state-level regression:

$$LFP_{i,t} = \alpha + \beta_1 t$$

where $LFP_{i,t}$ is prime-age labor-force participation for state i at date t (monthly).

Using the estimated coefficient, we predict the prime-age LFP for each state for March 2020 to December 2022, $\widehat{LFP}_{i,t}$. Figure A2 shows the actual (red line), as well as the estimated and predicted (green line) prime-age LFP for the US, Michigan (a state where LFP stays below trend during the sample), and Virginia (where LFP recovers rather quickly), as examples.

Figure A2: Estimated and Predicted LFP for the US, Michigan, and Virginia



Notes: Actual (red line) and predicted (green line) prime-age LFP. The figure shows the results for the entire United States and the states of Michigan and Virginia. Source: CDC


Then we obtain the difference as $Diff_{i,t} = LFP_{i,t} - \widehat{LFP}_{i,t}$. Negative values imply a lower labor-force participation than what has been predicted. To obtain the measure presented on the vertical axis of Figure 3, we average the difference for the period of July to December

2022.

A.3 Labor-Force Participation and Opioids

Figure A3 shows the correlation between the average prime-age LFP between 2017 and 2019, and alternative measures of opioid exposure: the percentage of individuals with opioid use disorder in 2017-2018 on the left, and the MME per capita of shipment of medications used to treat opioid addiction on the right. In both cases, the correlations are not statistically significant.

Figure A3: Relationship Between Prime-Age LFP in 2017-2019 and Alternative Opioid Exposure Measures




Notes: *Correlation between the average prime-age LFP between 2017 and 2019, and alternative measures of opioid exposure. Left panel: percentage of individuals with opioid use disorder in 2017-2018. Right panel: the MME per capita of shipment of medications used to treat opioid addiction. Source: CPS, NSDUH, ARCOS.*

Figure A4 shows the correlation between the growth in prime-age LFP between 2015 and 2019, and alternative measures of opioid exposure: the percentage of individuals with opioid use disorder in 2017-2018 on the left, and the MME per capita of shipment of medications used to treat opioid addiction on the right. In both cases, there is a positive correlation.

A.4 Stringency and Economic Support Index


Figure A5 shows the average and standard deviation of the Stringency and Economic Support index developed by the Oxford Tracker. The Oxford Tracker Dataset categorizes various COVID-19 policies into indexes representing different policy strengths, normalized between 0 and 100.

Figure A4: Relationship Between Prime-Age LFP Growth 2017-2019 and Alternative Opioid Exposure Measures



Notes: *Correlation between the prime-age LFP growth between 2015 and 2019, and alternative measures of opioid exposure. Left panel: percentage of individuals with opioid use disorder in 2017-2018. Right panel: the MME per capita of shipment of medications used to treat opioid addiction. Source: CPS, NSDUH, ARCOS.*

Figure A5: Stringency and Economic Support Index




Notes: *Summary policy indexes for Stringency and Economic Support reported by the Oxford Tracker for the entire United States and the standard deviation of these policy indexes across states. Source: Oxford Tracker.*

B Placebo, Heterogeneity and Robustness

B.1 Opioid Exposure and Opioid Deaths

Figure B1 shows the results obtained by estimating equation (1) using the opioid death rate as the dependent variable. The top panel uses the monthly opioid death rate, and the bottom panel uses the yearly age-adjusted opioid death rate. Initial opioid exposure is measured as a binary variable, with states being given a value of 1 if their age-adjusted opioid death rate in 2017 was above the median and 0 if below.

Figure B1: Non-Parametric Binary Event Study - Opioid Deaths




Notes: Non-parametric event study coefficients β_T and the 95% confidence interval. Coefficients represent changes in opioid overdose deaths relative to February 2020 between states with different levels of prior opioid exposure. The top panel uses the monthly crude rate, while the bottom panel uses the yearly age-adjusted rate. The initial opioid exposure measure has been dichotomized: states with an age-adjusted opioid death rate in 2017 above the median are given a value of 1, the others a value of 0.

B.2 Opioid Exposure and Drug Overdose Deaths

Figure B2 shows the results obtained by estimating equation (1) using drug overdose death rate as the dependent variable. The top panel uses the monthly death rate, and the bottom uses the yearly age-adjusted death rate. Initial opioid exposure is measured as a binary variable, with states being given a value of 1 if their age-adjusted opioid death rate in 2017 was above the median and 0 if below. The results are very similar to the ones obtained in Figure 5 in Section 4.

Figure B2: Non-Parametric Event Study with Continuous Exposure - Drug Overdose Deaths




Notes: Non-parametric event study coefficients β_T and the 95% confidence interval. Coefficients represent changes in drug overdose deaths relative to February 2020 between states with different levels of prior opioid exposure. The top panel uses the monthly crude rate, while the bottom uses the yearly age-adjusted rate. The initial opioid exposure measure has been normalized to have a standard deviation equal to 1.

B.3 Opioid Exposure and COVID-19 Cases

Figure B3 highlights that COVID-19 cases across states were uncorrelated with pre-COVID opioid exposure. The figure shows the estimates obtained by running the non-parametric event study, equation (1), using monthly COVID-19 cases per 100,000 people as the dependent variable, and the continuous measure of initial opioid exposure as the

independent variable. The analysis is done only for the post-COVID period.

Figure B3: Non-Parametric Event Study - COVID-19 Cases



Notes: *Non-parametric event-study coefficients β_T and the 95% confidence interval. Coefficients represent changes in total COVID-19 cases reported relative to February 2020 between states with different levels of prior opioid exposure. The initial opioid exposure measure has been normalized to have a standard deviation equal to 1.*


B.4 Placebo Analysis

Table B1 shows the age-adjusted death rate in 2017 across the 15 leading causes of death in the US. The initial exposure measure used for the placebo analysis in the main text is constructed by aggregating the state-level death rate from these 15 causes, excluding the third (accidents) and eighth (intentional self-harm) causes of death. The reason for this is that some of the deaths in these causes are also included in the opioid death rate measure. Figure B4 presents the estimated coefficients from the parametric event study where the initial exposure measure is based on the total death rate from only the top 10 causes, excluding the third and eighth. Finally, Figure B5 shows the estimated coefficients from parametric event studies that have as the initial exposure measure the death rate of the cause highlighted in the title.

Table B1: Average Age-Adjusted Death Rate by Cause in 2017


	Death Rate
1. Diseases of heart (I00-I09,I11,I13,I20-I51)	165.99
2. Malignant neoplasms (C00-C97)	155.00
3. Accidents (unintentional injuries) (V01-X59,Y85-Y86)	54.01
4. Chronic lower respiratory diseases (J40-J47)	43.78
5. Cerebrovascular diseases (I60-I69)	37.43
6. Alzheimer disease (G30)	32.07
7. Diabetes mellitus (E10-E14)	21.99
8. Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	16.50
9. Influenza and pneumonia (J09-J18)	14.90
10. Nephritis (N00-N07,N17-N19,N25-N27)	12.79
11. Chronic liver disease and cirrhosis (K70,K73-K74)	11.30
12. Septicemia (A40-A41)	10.27
13. Parkinson disease (G20-G21)	8.62
14. Essential hypertension and hypertensive renal disease (I10,I12,I15)	8.44
15. Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6.69

Notes: The table shows the age-adjusted death rate in 2017 by the leading causes of death in the US. Source: CDC - Underlying Cause of Death.

Figure B4: Parametric Event Study with Placebo Exposure

Notes: Placebo parametric event study coefficients θ_T and the 95% confidence interval when using prime-age LFP as the dependent variable. Coefficients represent average differences from state-trend in prime-age LFP between states with different levels of prior placebo exposure. The measure of prior placebo exposure is the total age-adjusted death rate in 2017 due to the top 10 underlying causes of death. The death rate has been normalized to have a standard deviation equal to 1.

Figure B5: Parametric Event Study Estimates - Placebo




Notes: Placebo parametric event study coefficients θ_T and the 95% confidence interval. Coefficients represent average differences from state-trend in prime-age LFP between states with different levels of prior placebo exposure. The measure of prior placebo exposure is the age-adjusted death rate in 2017 due to causes indicated in the title of each panel. All death rates have been normalized to have a standard deviation equal to 1.

B.5 Alternative Age Groups

The main analysis focuses on the labor-force participation of prime-age workers, i.e., ages between 25 and 54. Figure B6 shows the estimated coefficients from the parametric event study analysis when focusing on the age groups 18-24 and 55-64.

Figure B6: Parametric Event Study Estimates - Labor Force Across Samples




Notes: *Parametric event study coefficients θ_T and the 95% confidence interval. Coefficients represent average differences from state-trend in prime-age LFP between states with different levels of prior opioid exposure. The upper panel restricts the sample to individuals aged 18-24, and the lower panel to those 55-64. The initial opioid exposure measure has been normalized to have a standard deviation equal to 1.*

B.6 Alternative Exposure Measures

Figure B7 shows the estimated coefficients from the parametric event study analysis when using two alternative measures of opioid exposure are presented: the logarithm of the age-adjusted opioid death rate in 2017 and the average age-adjusted opioid death rate between 2010 and 2017.

Figure B7: Parametric Event Study Estimates - Measures of Opioid Death Rate



Notes: *Parametric event study coefficients θ_T and the 95% confidence interval. Coefficients represent average differences from state-trend in prime-age LFP between states with different levels of prior opioid exposure. Prior opioid exposure in the top panel is given by the log of the opioid-related overdose age-adjusted death rate. The bottom panel shows the average opioid-related overdose age-adjusted death rate between 2010 and 2017, normalized to have a standard deviation equal to 1.*

C Data on Mechanisms

C.1 Employment and Health Statistics Across Opioid Users

Tables C1 and C2 present data on employment, health, and work disability for adults aged 24–49, based on the 2019 and 2021/2022 waves of the NSDUH. Individuals are categorized by their drug use in the past 12 months. The nonuser group includes those who reported no drug use, excluding marijuana. The disorder group comprises individuals diagnosed with a drug use disorder, excluding marijuana use disorder.

Table C1: Employment Status by Drug Use, Ages 24-49

<i>Panel A</i>		2019		2021/2022		
	Emp.	Unemp.	Out of LF	Emp.	Unemp.	Out of LF
Nonuser	79.3%	3.9%	16.8%	75.8%	5.1%	19.1%
Disorder	58.7%	15.3%	26.0%	54.3%	13.0%	32.7%
Population	79.0%	4.4%	16.6%	75.0%	5.7%	19.3%

<i>Panel B</i>		2019		2021/2022		
	Very Good	Good	Fair/Poor	Very Good	Good	Fair/Poor
Nonuser	61.4%	28.7%	9.9%	58.1%	31.2%	10.7%
Disorder	39.5%	35.0%	25.5%	33.2%	37.4%	29.4%
Population	60.5%	28.9%	10.6%	56.5%	31.7%	11.8%

Notes: The table shows the employment status in Panel A, and self-reported health status in Panel B, by type of drug user in 2019 and for the 2021/2022 period. Self-reported health status is a categorical variable with 4 options: very good, good, fair, and poor. Fair and poor categories are aggregated. The type of drug user is defined by the NSDUH. See the text for further details. Nonusers do not use any drug, excluding marijuana. For each drug user type and year, the row sums up to 100. Source: NSDUH.


C.2 Out-of-labor-force due to Disability Rate

Figure C1 shows the share of prime-age individuals who do not work due to disability.

Table C2: Drug Use and Work Disability, Ages 24-49

	NLF - Disability		Work Days Sick		Work Days Skip	
	2019	2021/2022	2019	2021/2022	2019	2021/2022
Nonuser	17.4%	16.8%	0.83	1.00	0.28	0.32
Disorder	25.1%	44.1%	2.12	2.08	1.94	0.96
Population	18.2%	19.4%	0.88	1.05	0.34	0.38

Notes: The table shows several measures of absence from work by type of opioid use in 2019 and for the 2021/2022 period. The first two columns show the percentage of people out of the labor force due to a disability. The second two columns show the average number of work days missed in the past 30 days due to sickness. The last two columns show the average number of work days missed for other reasons. The type of opioid use is defined by the NS-*DUH*. See the text for further details. Source: NS*DUH*


Figure C1: Out-of-labor-force due to Disability Rate

Notes: Share of prime-age individuals not in the labor force for disability reasons. This measure includes individuals who report being unable to work and not working due to disability. Source: CPS

C.3 Out-of-labor-force due to Disability Rate across Education

Figure C2 shows the estimated coefficients from conducting separate parametric event studies for the non-college and college-educated samples using the fraction out-of-labor-force due to disability as the dependent variable.

Figure C2: Parametric Event Study Estimates - Out-of-LF due to Disability



Notes: *Parametric event-study coefficients θ_T and the 95% confidence interval. Coefficients represent average differences from state-trend in prime-age out of labor force due to disability between states with different levels of prior opioid exposure. The top panel uses prime-age out of labor force due to disability among non-college-educated, while the bottom uses college-educated. The initial opioid exposure measure has been normalized to have a standard deviation equal to 1.*